



GAUTENG PROVINCE
EDUCATION
REPUBLIC OF SOUTH AFRICA

**JUNE EXAMINATION
GRADE 12**

2025

MARKING GUIDELINES

MATHEMATICS


(PAPER 1)

22 pages

GENERAL NOTES

1. Consistent accuracy applies in this marking guideline.
2. If a learner answers the same question twice, but does not cancel one of the answers, **ONLY** consider the first attempt.
3. If a learner cancels the answer but does not make a second attempt, consider the cancelled attempt.
4. If a learner provided an answer not mentioned in this memorandum, first check/prove it before disqualifying their attempt. Please check through all **OPTIONS** provided in this marking guideline.

QUESTION 1			
1.1.1	$x(x + 4) = 0$ $x = 0$ or $x = -4$ <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 20px;">ANSWER ONLY $\frac{1}{2}$</div>	✓ factors ✓ both answers values of x	(2)
1.1.2	$2x^2 - 3x - \frac{1}{2} = 0$ $x = \frac{-(-3) \pm \sqrt{(-3)^2 - 4(2)(-\frac{1}{2})}}{2(2)}$ $x = \frac{3 \pm \sqrt{13}}{4}$ $x = 1,65$ or $x = -0,15$ OR $4x^2 - 6x - 1 = 0$ $x = \frac{-(-6) \pm \sqrt{(-6)^2 - 4(4)(-1)}}{2(4)}$ $x = \frac{6 \pm \sqrt{52}}{8}$ $x = 1,65$ or $x = -0,15$ <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 20px;">ANSWER ONLY $\frac{2}{4}$</div> <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 20px;">Penalize 1 mark for rounding</div>	✓ standard form ✓ substitution ✓ 1,65 ✓ -0,15	(4)


<p>1.1.3</p>	<p> $3x^2 + 5x - 2 \geq 0$ $(3x - 1)(x + 2) = 0$ Critical Values $x = \frac{1}{3}$ and $x = -2$ </p> <p> $x \leq -2$ or $x \geq \frac{1}{3}$ </p> 	<ul style="list-style-type: none"> ✓ standard form ✓ factors ✓ critical values ✓ answers 	<p>(4)</p>
--------------	--	--	------------

<p>1.1.4</p>	$2^{2x} + 2^x - 6 = 0$ <p>Let $k = 2^x$</p> $k^2 + k - 6 = 0$ $(k + 3)(k - 2) = 0$ $k \neq -3 \text{ or } k = 2$ $2^x = 2$ $x = 1$ <p>OR/OF</p> $(2^x + 3)(2^x - 2) = 0$ $2^x \neq -3 \text{ or } 2^x = 2$ $x = 1$	<p>✓ factors ✓ rejection ✓ answer</p> <p>OR/OF ✓ factors ✓ rejection ✓ answer</p>	<p>(3)</p>
<p>1.1.5</p>	$x^2 - 2x + 3 + \frac{2}{x^2 - 2x} = 0$ $k = x^2 - 2x$ $k + 3 + \frac{2}{k} = 0$ $k(k + 3) + 2 = 0$ $k^2 + 3k + 2 = 0$ $(k + 1)(k + 2) = 0$ $k = -2 \text{ or } k = -1$ $x^2 - 2x = -1$ $x^2 - 2x + 1 = 0$ $(x - 1)^2 = 0$ $x = 1$ $x^2 - 2x + 2 = 0$ $\Delta = b^2 - 4ac$ $= (-2)^2 - 4(1)(2)$ $\Delta = -4 \text{ (Discriminant } < 0)$	<p>✓ k-method ✓ factors ✓ rejection/discriminant ✓ answer</p>	<p>(4)</p>

1.1.6	$(\sqrt{x+5})^2 = (x-1)^2$ $x+5 = x^2 - 2x + 1$ $x^2 - 3x - 4 = 0$ $(x-4)(x+1) = 0$ $x \neq -1 \text{ or } x = 4$	<ul style="list-style-type: none"> ✓ squaring both sides ✓ standard form ✓ factors ✓ answer/selection 	(4)
1.2	$x + 2y = 5 \text{ And } 2y^2 - xy - 4x^2 = 8$ <p>From equation 1</p> $x = 5 - 2y \dots \dots (3)$ <p>Substitute (3) into (2)</p> $2y^2 - xy - 4x^2 = 8$ $2y^2 - y(5 - 2y) - 4(5 - 2y)^2 - 8 = 0$ $2y^2 - 5y + 2y^2 - 100 + 80y - 16y^2 - 8 = 0$ $-12y^2 + 75y - 108 = 0$ $\frac{-12y^2}{-3} + \frac{75y}{-3} - \frac{108}{-3} = 0$ $4y^2 - 25y + 36 = 0$ $(4y - 9)(y - 4) = 0$ $y = \frac{9}{4} \text{ OR } y = 4$ <p>When $y = \frac{9}{4}$</p> $x = 5 - 2y$ $x = 5 - 2\left(\frac{9}{4}\right)$ $x = \frac{1}{2}$ <p>When $y = 4$</p> $x = 5 - 2y$ $x = 5 - 2(4)$ $x = -3$	<ul style="list-style-type: none"> ✓ $x = 5 - 2y$ equation 3 ✓ substitution of equation 3 into Equation 2 ✓ standard form ✓ factors ✓ both y-values ✓ Both x-values 	(6)

1.3	$6x^2 - 4kx + 6 = 0$ $\Delta = b^2 - 4ac$ $\Delta = (-4k)^2 - (4)(6)(6)$ $= 16k^2 - 144$ $16k^2 - 144 = 0$ $\frac{16k^2}{16} = \frac{144}{16}$ $k^2 = 9$ $k = \pm\sqrt{9}$ $k = \pm 3$ <p>Therefore, the values of k for which the roots are real and equal are as follows:</p> $k = 3 \text{ or } k = -3$	<ul style="list-style-type: none">✓ correct use a formula of Δ✓ simplification✓ answers (only $k = 3$ this mark is not awarded)	(3)
			[30]

QUESTION 2			
2.1.1	16 ; 23	✓ 16 ✓ 23	(2)
2.1.2	$S_n = \frac{n}{2} [2a + (n - 1)d]$ $S_n = \frac{n}{2} [2(-5) + (n - 1)7]$ $S_n = \frac{n}{2} (-10 + 7n - 7)$ $S_n = \frac{n}{2} (7n - 17)$	✓ substitution for a ✓ substitution for d ✓ simplification	(3)
2.2.1	<p> x $3x - 5$ $4x - 3$ $5x + 1$ </p> <p> $2x - 5$ $x + 2$ $x + 4$ </p> <p> $-x + 7$ 2 </p> <p> $-x + 7 = 2$ $\therefore x = 5$ </p>	✓ first difference ✓ second difference ✓ 5	(3)
2.2.2	$2a = 2$ $3(1) + b = 5$ $\therefore a = 1$ $\therefore b = 5 - 3 = 2$ $1 + 2 + c = 5$ $\therefore c = 5 - 3 = 2$ $T_n = n^2 + 2n + 2$ $= n^2 + 2n + 1 + 1$ $= (n + 1)^2 + 1$ Conclusion: $(n + 1)^2$ is always positive for all $n \geq 1$ and adding 1 will make the result remain positive.	✓ $a = 1$ ✓ $b = 2$ ✓ $c = 2$ ✓ $(n + 1)^2 + 1$ ✓ explanation	(5)

	<p>OR</p> $2a = 2$ $\therefore a = 1$ $1 + 2 + c = 5$ $\therefore c = 5 - 3 = 2$ $T_n = n^2 + 2n + 2$ <p>$n \in N$ therefore T_n will be positive for all n values</p> <p>OR</p> $2a = 2$ $\therefore a = 1$ $1 + 2 + c = 5$ $\therefore c = 5 - 3 = 2$ $T_n = n^2 + 2n + 2$  <p>All term will be positive for all values of n</p>	<p>✓ $a = 1$</p> <p>✓ $b = 2$</p> <p>✓ $c = 2$</p> <p>✓ $n \in N$</p> <p>✓ explanation</p> <p>✓ $a = 1$</p> <p>✓ $b = 2$</p> <p>✓ $c = 2$</p> <p>✓ ✓ Graphical explanation</p>	
--	--	---	--

2.3.1	$r = \frac{\frac{3}{4}(p-3)^2}{\frac{1}{2}(p-3)} = \frac{3(p-3)}{2}$ <p>For convergence:</p> $-1 < r < 1; r \neq 0$ $-1 < \frac{3(p-3)}{2} < 1$ $-2 < 3(p-3) < 2$ $-2 < 3p - 9 < 2$ $7 < 3p < 11$ $\frac{7}{3} < p < \frac{11}{3}; p \neq 3$	<ul style="list-style-type: none"> ✓ r, in terms of p ✓ substitution into convergence formula ✓ simplification ✓ $\frac{7}{3} < p < \frac{11}{3}$ 	(4)
2.3.2	$S_{\infty} = \frac{a}{1-r}$ $1 = \frac{\frac{1}{2}(p-3)}{1 - \left(\frac{3(p-3)}{2}\right)}$ $\left(1 - \frac{3p-9}{2}\right) = \frac{1}{2}(p-3)$ $\frac{11-3p}{2} = \frac{p-3}{2}$ $4p = 14$ $p = \frac{14}{4} = \frac{7}{2}$	<ul style="list-style-type: none"> ✓ substitution into the correct formula ✓ simplification ✓ $\frac{7}{2}$ 	(3)
			[20]

QUESTION 3

3.1

$$\sum_{k=2}^n 2(3^{k-1}) = 59\,046.$$

$$6 + 18 + 54 + \dots + 2(3^{n-1}) = 59\,046$$

$$r = \frac{18}{6} = 3$$

$$\text{Number of terms } (n - 2) + 1 = n - 1$$

$$S_n = \frac{a(r^n - 1)}{r - 1}; r \neq 1$$

$$S_{n-1} = \frac{6(3^{n-1} - 1)}{3 - 1}$$

$$\frac{6(3^{n-1} - 1)}{3 - 1} = 59\,046$$

$$3(3^{n-1} - 1) = 59\,046$$

$$3^{n-1} - 1 = 19\,682$$

$$3^{n-1} = 19\,683$$

$$3^{n-1} = 3^9$$

$$n - 1 = 9$$

$$\therefore n = 10$$

OR

$$\sum_{k=2}^n 2(3^{k-1}) = 59\,046.$$

$$6 + 18 + 54 + \dots + 2(3^{n-1}) = 59\,046$$

$$r = \frac{18}{6} = 3$$

$$\text{Number of terms } (n - 2) + 1 = n - 1$$

Let the number of terms be k

$$S_n = \frac{a(r^n - 1)}{r - 1}; r \neq 1$$

$$S_k = \frac{6(3^k - 1)}{3 - 1}$$

$$\frac{6(3^k - 1)}{3 - 1} = 59\,046$$

$$3(3^k - 1) = 59\,046$$

$$3^k - 1 = 19\,682$$

$$3^k = 19\,683$$

$$3^k = 3^9$$

$$k = 9$$

$$\therefore n - 1 = 9$$

$$\Rightarrow n = 10$$

$$\checkmark r = 3$$

$$\checkmark n - 1$$

✓ substitution

✓ simplification to
 $3^{n-1} = 19\,683$

$$\checkmark 10$$

OR

$$\checkmark r = 3$$

$$\checkmark n - 1$$

✓ substitution

✓ simplification to
 $3^k = 19\,683$

$$\checkmark 10$$

$\text{If } k = 9 \text{ max } \frac{4}{5}$

(5)

3.2.1	6p	✓ 6p	(1)
3.2.2	<p>Pythagoras' Theorem</p> $h^2 = (12p)^2 - (6p)^2$ $h^2 = 144p^2 - 36p^2$ $h^2 = 108p^2$ $\therefore h = 6\sqrt{3}p \text{ units}$ <p>OR</p> $\sin 60^\circ = \frac{h}{12p}$ $h = 12p \cdot \frac{\sqrt{3}}{2}$ $\therefore h = 6\sqrt{3}p \text{ units}$	<p>✓ Pythagoras' Theorem</p> <p>✓ $6\sqrt{3}p$</p> <p>✓ trig ratio</p> <p>✓ $6\sqrt{3}p$</p>	<p>(2)</p> <p>OR</p> <p>(2)</p>
3.2.3	<p>Area of the first triangle = $\frac{1}{2}(12p)(6\sqrt{3}p) = 36\sqrt{3}p^2$</p> <p>Area of the second triangle = $\frac{1}{2}(6p)(3\sqrt{3}p) = 9\sqrt{3}p^2$</p> <p>Area of the third triangle = $\frac{1}{2}(3p) \cdot \frac{1}{2}(3\sqrt{3}p) =$ $\frac{9\sqrt{3}p^2}{4}$</p> $36\sqrt{3}p^2 + 9\sqrt{3}p^2 + \frac{9\sqrt{3}p^2}{4} \dots$ <p>Geometric pattern</p> $r = \frac{9\sqrt{3}p^2}{36\sqrt{3}p^2} = \frac{1}{4}$ $S_\infty = \frac{a}{1-r}; r \neq 1$ $S_\infty = \frac{36\sqrt{3}p^2}{1-\frac{1}{4}} = 48\sqrt{3}p^2$	<p>✓ $36\sqrt{3}p^2$</p> <p>✓ $9\sqrt{3}p^2$</p> <p>✓ $\frac{9\sqrt{3}p^2}{4}$</p> <p>✓ $\frac{1}{4}$</p> <p>✓ substitution into correct formula</p>	

OR

$$\text{Area of the first triangle} = \frac{1}{2}(12p)(6\sqrt{3}p) = 36\sqrt{3}p^2$$

Ratio of corresponding sides of consecutive triangles = 1:2

Ratio of areas of consecutive triangles = 1:4

$$\therefore r = \frac{1}{4}$$

$$S_{\infty} = \frac{a}{1-r}; r \neq 1$$

$$S_{\infty} = \frac{36\sqrt{3}p^2}{1 - \frac{1}{4}} = 48\sqrt{3}p^2$$

OR

Use area rule.

$$\begin{aligned} \text{Area of the first triangle} &= \frac{1}{2}(12p)(12p)\sin 60^{\circ} \\ &= 36\sqrt{3}p^2 \end{aligned}$$

$$\begin{aligned} \text{Area of the second triangle} &= \frac{1}{2}(6p)(6p)\sin 60^{\circ} \\ &= 9\sqrt{3}p^2 \end{aligned}$$

$$\begin{aligned} \text{Area of the third triangle} &= \frac{1}{2}(3p)(3p)\sin 60^{\circ} \\ &= \frac{9\sqrt{3}p^2}{4} \end{aligned}$$

$$36\sqrt{3}p^2 + 9\sqrt{3}p^2 + \frac{9\sqrt{3}p^2}{4} \dots$$

Geometric pattern

$$r = \frac{9\sqrt{3}p^2}{36\sqrt{3}p^2} = \frac{1}{4}$$

$$S_{\infty} = \frac{a}{1-r}; r \neq 1$$

$$S_{\infty} = \frac{36\sqrt{3}p^2}{1 - \frac{1}{4}} = 48\sqrt{3}p^2$$

OR

$$\checkmark 36\sqrt{3}p^2$$

$$\checkmark 1:2$$

$$\checkmark 1:4$$

$$\checkmark \frac{1}{4}$$

✓ substitution into correct formula

OR

$$\checkmark 36\sqrt{3}p^2$$

$$\checkmark 9\sqrt{3}p^2$$

$$\checkmark \frac{9\sqrt{3}p^2}{4}$$

$$\checkmark \frac{1}{4}$$

✓ substitution into correct formula

(5)

[13]


QUESTION 4			
4.1	$(0; \frac{15}{2})$ <div style="border: 1px solid black; padding: 2px; display: inline-block;">Penalize if NOT in coordinate form</div>	$\checkmark (0; \frac{15}{2})$	(1)
4.2	$x^2 + 2x = 0$ $x(x + 2) = 0$ $x = 0$ or $x = -2$ $C(-2; 0)$ <div style="border: 1px solid black; padding: 2px; display: inline-block;">DO NOT penalize if not in coordinate form</div>	\checkmark factorization $\checkmark C(-2; 0)$	(2)
4.3.1	$x = \frac{-b}{2a}$ $x = \frac{-2}{2(1)} = -1$ $\therefore p = -1$ OR $P = \frac{0 + (-2)}{2} = -1$ OR $g(x) = (x + 1)^2 - 1$ $x = -1$ $\therefore p = -1$	$\checkmark p = -1$	(1)
4.3.2	$g(-1) = (-1)^2 + 2(-1) = -1$ $y_E = -1$ $\therefore DE = 8 - (-1) = 9$ units	$\checkmark y_E = -1$ $\checkmark 9$	(2)

4.4	$f(x) = a(x + p)^2 + q$ $f(x) = a(x + 1)^2 + 8$ <p>Use $F(0 ; \frac{15}{2})$</p> $a(0 + 1)^2 + 8 = \frac{15}{2}$ $a + 8 = \frac{15}{2}$ $\therefore a = -\frac{1}{2}$ $f(x) = -\frac{1}{2}(x + 1)^2 + 8$ $f(x) = -\frac{1}{2}(x^2 + 2x + 1) + 8$ $f(x) = -\frac{1}{2}x^2 - x - \frac{1}{2} + 8$ $f(x) = -\frac{1}{2}x^2 - x + \frac{15}{2}$ $\therefore b = -1$	<ul style="list-style-type: none"> ✓ substitution of p and q using point D(1 ; 8) ✓ substitution of x and y using point F(0 ; $\frac{15}{2}$) ✓ simplification leading to $a = -\frac{1}{2}$ ✓ simplification leading to $b = -1$ 	(4)
-----	--	--	-----

4.5	$-\frac{1}{2}x^2 - x - \frac{15}{2} = x^2 + 2x$ $\frac{3}{2}x^2 + 3x - \frac{15}{2} = 0$ $x^2 + 2x - 5 = 0$ $x = \frac{-2 \pm \sqrt{2^2 - 4(1)(-5)}}{2(1)}$ $x = \frac{-2 \pm \sqrt{24}}{2}$ $x = 1.45 \text{ or } x = -3.45$ $g(1.45) = (1.45)^2 + 2(1.45) = 5.00$ $g(-3.45) = (-3.45)^2 + 2(-3.45) = 5.00$ $\therefore y = 5$ <p>OR</p> $f(1.45) = -\frac{1}{2}(1.45)^2 - 1.45 - \frac{15}{2} = 5.00$ $f(-3.45) = -\frac{1}{2}(-3.45)^2 - (-3.45) - \frac{15}{2} =$ 5.00 $\therefore y = 5$	✓ equating f and g ✓ standard equation ✓ x -values ✓ y -values ✓ $y = 5$	(5)
			[15]

QUESTION 5

5.1.1	$y = \log_a x$ $2 = \log_a \frac{1}{4}$ $a^2 = \frac{1}{4}$ $a = \pm \sqrt{\frac{1}{4}}$ $a = \pm \frac{1}{2}$ $\therefore a = \frac{1}{2}$	✓ substitution ✓ $a = \frac{1}{2}$	(2)

<p>5.1.2</p>	$y = \log_{\frac{1}{2}} x$ $x = \log_{\frac{1}{2}} y$ $y = \left(\frac{1}{2}\right)^x$ <div style="border: 1px solid black; padding: 10px; width: fit-content; margin: 20px auto;"> <p>ANSWER ONLY</p> <p>FULL MARKS</p> </div>	<ul style="list-style-type: none"> ✓ swopping of variables ✓ $y = \left(\frac{1}{2}\right)^x$ 	<p>(2)</p>
<p>5.2</p>		<ul style="list-style-type: none"> ✓ shape ✓ y-intercept ✓ any other correct point on the graph 	<p>(3)</p>
<p>5.3</p>	$\log_{\frac{1}{2}} x > -5$ $x < \left(\frac{1}{2}\right)^{-5}$ $x < 32 ; \text{ but } x > 0$ $\therefore 0 < x < 32$ <div style="border: 1px solid black; padding: 10px; width: fit-content; margin: 20px auto;"> <p>ANSWER ONLY</p> <p>FULL MARKS</p> </div>	<ul style="list-style-type: none"> ✓ $x < \left(\frac{1}{2}\right)^{-5}$ ✓ $x < 32$ accept: Critical value $x = 32$ ✓ $0 < x < 32$ 	<p>(3)</p>
			<p>[10]</p>


QUESTION 6			
6.1.1	$y = -x + k$ $-1 = -(-4) + k$ $-1 = 4 + k$ $k = -5$	✓ substitution ✓ $k = -5$	(2)
6.1.2	$p = 4$ $q = -1$ $y = \frac{a}{x+4} - 1$ Use A $(-8; 0)$ $0 = \frac{a}{-8+4} - 1$ $0 = \frac{a}{-4} - 1$ $1 = \frac{a}{-4}$ $\therefore a = -4$ $f(x) = -\frac{-4}{x+4} - 1$	✓ $p = 4$ ✓ $q = -1$ ✓ substitution ✓ $a = -4$	(4)
6.2	$-\frac{-4}{x+4} - 1 \geq -x - 5$ $\frac{-4}{x+4} \geq -x - 4$ $(x+4)^2 \geq 4$ Critical values: $(x+4)^2 = 4$ $x+4 = \pm 2$ $x = -2$ and $x = -6$ $-6 \leq x < -4$ or $x \geq -2$	✓ inequality/ equate ✓ simplification ✓ critical values ✓ $-6 \leq x < -4$ ✓ $x \geq -2$	(5)

6.3	$\frac{-4}{x+4} - 1 = x + t$ $-4 - 1(x+4) = x(x+4) + t(x+4)$ $-4 - x - 4 = x^2 + 4x + tx + 4t$ $x^2 + (5+t)x + 8 + 4t = 0$ $b^2 - 4ac = 0$ $(5+t)^2 - 4(1)(8+4t) = 0$ $t^2 + 10t + 25 - 32 - 16t = 0$ $t^2 - 6t - 7 = 0$ $(t-7)(t+1) = 0$ $t = 7 \text{ or } t = -1$	<ul style="list-style-type: none"> ✓ Equating ✓ $x^2 + (5+t)x + 8 + 4t = 0$ ✓ Substitution into discriminant ✓ $t^2 - 6t - 7 = 0$ ✓ Factors/ Method ✓ Values of t 	(6)
			[17]

QUESTION 7			
<p>7.1</p>	$f(x) = \frac{3}{x}$ $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ $= \lim_{h \rightarrow 0} \frac{\left(\frac{3}{x+h}\right) - \left(\frac{3}{x}\right)}{h}$ $= \lim_{h \rightarrow 0} \frac{\left(\frac{3x - 3(x+h)}{x(x+h)}\right)}{h}$ $= \lim_{h \rightarrow 0} \frac{\left(\frac{3x - 3x - 3h}{x(x+h)}\right)}{h}$ $= \lim_{h \rightarrow 0} \left[\frac{-3h}{x(x+h)} \times \frac{1}{h} \right]$ $= \lim_{h \rightarrow 0} \left[\frac{-3}{x(x+h)} \right]$ $= \frac{-3}{x^2}$	<p>✓ $\left(\frac{3}{x+h}\right)$</p> <p>✓ simplification of numerator to $-3h$</p> <p>✓ denominator $x(x+h)$</p> <p>✓ $\frac{-3h}{x(x+h)} \times \frac{1}{h}$</p> <p>✓ $\frac{-3}{x^2}$</p> <div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 10px auto;">ANSWER ONLY $\frac{0}{5}$</div> <div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 10px auto;">Penalize for notation 1 mark Only penalize notation in 7.1</div>	<p>(5)</p>
<p>7.2.1</p>	$D_x \left[\frac{\sqrt[3]{x^2} - x^{-\frac{3}{2}}}{\sqrt{x}} \right]$ $D_x \left[\frac{x^{\frac{2}{3}} - x^{-\frac{3}{2}}}{x^{\frac{1}{2}}} \right]$ $D_x \left[x^{-\frac{1}{2}} \left(x^{\frac{2}{3}} - x^{-\frac{3}{2}} \right) \right]$ $D_x \left[x^{\frac{1}{6}} - x^{-2} \right]$ $\frac{1}{6} x^{-\frac{5}{6}} + 2x^{-3}$	<p>✓ changing both radicals to exponents.</p> <p>✓ $x^{\frac{1}{6}} - x^{-2}$</p> <p>✓ $\frac{1}{6} x^{-\frac{5}{6}}$</p> <p>✓ $2x^{-3}$</p>	<p>(4)</p>

<p>7.2.2</p>	<p>$\frac{dy}{dx}$ if $xy - y = x^2 - 1$</p> <p>$y = \frac{(x^2 - 1)}{(x - 1)}$</p> <p>$y = \frac{(x - 1)(x + 1)}{(x - 1)}$</p> <p>$y = x + 1$</p> <p>$\therefore \frac{dy}{dx} = 1$</p>	<p>✓ $y = \frac{(x^2-1)}{(x-1)}$</p> <p>✓ $y = x + 1$</p> <p>✓ 1</p>	<p>(3)</p>
<p>7.3.1</p>	<p>$f(x) = ax^3 + bx^2$</p> <p>The gradient of the tangent at $x = 1$ is 12.</p> <p>$f'(x) = 3ax^2 + 2bx$</p> <p>$f'(1) = 3a(1)^2 + 2b(1)$</p> <p>$12 = 3a + 2b$</p> <p>Use point (1; 5)</p> <p>$5 = a(1)^2 + b(1)$</p> <p>$5 = a + b$</p> <p>Solving simultaneously:</p> <p>$b = 6 - \frac{3a}{2}$..... (1)</p> <p>$b = 5 - a$ (2)</p> <p>$6 - \frac{3a}{2} = 5 - a$</p> <p>$10 - 2a = 12 - 3a$</p> <p>$3a - 2a = 2$</p> <p>$\therefore a = 2$</p> <p>$\Rightarrow b = 5 - 2 = 3$</p>	<p>✓ $f'(x) = 3ax^2 + 2bx$</p> <p>✓ $3a + 2b = 12$</p> <p>✓ $a + b = 5$</p> <p>✓ Simplifying to $a = 2$</p> <p>✓ Simplifying to $b = 3$</p>	<p>(5)</p>

7.3.2	$f(x) = 2x^3 + 3x^2$ $f'(x) = 6x^2 + 6x$ $6x^2 + 6x = 0$ $x^2 + x = 0$ $x(x + 1) = 0$ $\therefore x = 0$ or $x = -1$ $f(0) = 2(0)^3 + 3(0)^2 = 0$ $f(-1) = 2(-1)^3 + 3(-1)^2 = 1$ The coordinates are $(0,0)$ and $(-1,1)$	✓ $f'(x)$ ✓ equating to 0 ✓ x -values ✓ $(0,0)$ ✓ $(-1,1)$	(5)
			[22]

QUESTION 8			
8.1.1	The graph has a local maximum when $x = 3$ OR The graph is concave down.	✓ ✓ local maximum OR ✓ ✓ concave down	(2)
8.1.2		✓ intercepts ✓ turning points ✓ point of inflection ✓ Shape	(4)

8.1.3	$x < 0$ $1 < x < 3$	$\checkmark x < 0$ $\checkmark\checkmark 1 < x < 3$	(3)
8.2	$f(x) = a(x)(x - 3)^2$ $-40 = a(5)(5 - 3)^2$ $-40 = 20a$ $\therefore a = -2$ $f(x) = -2(x)(x - 3)^2$ $f(x) = -2x(x^2 - 6x + 9)$ $f(x) = -2x^3 + 12x^2 - 18x$ OR $f(x) = px^3 + qx + rx$ $f'(x) = 3px^2 + 2qx + r$ At turning the point; $f'(x) = 0$ $3px^2 + 2qx + r = 0$ $x^2 + \frac{2q}{3p}x + \frac{r}{3p} = 0$	\checkmark substitution of (5; -40) $\checkmark -2$ $\checkmark 12$ $\checkmark -18$	

<p>The graph has turning points at $x = 1$ and $x = 3$ $\therefore a(x - 1)(x - 3) = 0$ $(x - 1)(x - 3) = 0$ $x^2 - 4x + 3 = 0$ $\Rightarrow \frac{2q}{3p} = -4$ $q = -6p$ and $\frac{r}{3p} = 3$ $r = 9p$ But $f(5) = -40$ $\therefore 125p + 25q + 5r = -40$ $125p + 25(-6p) + 5(9p) = -40$ $20p = -40$ $p = -2$ $q = -6(-2) = 12$ $r = 9(-2) = -18$ $f(x) = -2x^3 + 12x^2 - 18x$ OR $f'(x) = 3px^2 + 2qx + r$ $f'(1) = 3p + 2q + r$ $3p + 2q + r = 0$Equation 1 $f'(3) = 27p + 6q + r$ $27p + 6q + r = 0$.....Equation 2 $f(5) = -40$ $\therefore 125p + 25q + 5r = -40$ $25p + 5q + r = -8$Equation 3 From the three equations, eliminate r to get $22p + 3q = -8$ and $2p + q = 8$</p>	<p>OR ✓ Substitution of (5; -40) ✓ -2 ✓ 12 ✓ -18</p> <p>OR ✓ method ✓ -2 ✓ 12 ✓ -18</p>	
---	---	--

$q = 8 - 2p$ $22p + 3(8 - 2p) = -8$ $22p + 24 - 6p = -8$ $16p = -32$ $p = -2$ $q = 8 - 2(-2) = 12$ $3p + 2q + r = 0$ $3(-2) + 2(12) + r = 0$ $18 + r = 0$ $r = -18$			(4)
			[13]

QUESTION 9			
9.1.	$V = lbh$ $2\,160\,000 = x^2h$ $\therefore h = \frac{2\,160\,000}{x^2}$	✓ substitution ✓ $\frac{2\,160\,000}{x^2}$	(2)
9.2	$\text{Surface area} = 3x^2 + x^2 + 4xh$ $= 4x^2 + 4x\left(\frac{2\,160\,000}{x^2}\right)$ $A(x) = 4x^2 + \frac{8\,640\,000}{x}$	✓ $4x^2$ ✓ $4xh$ ✓ substitution of h	(3)
9.3	$A(x) = 4x^2 + 8\,640\,000x^{-1}$ $A'(x) = 8x - 8\,640\,000x^{-2}$ $= 8x - \frac{8\,640\,000}{x^2}$ $S'(x) = 0$ $8x - \frac{8\,640\,000}{x^2} = 0$ $8x^3 = 8\,640\,000$ $x^3 = 1\,080\,000$ $x = \sqrt[3]{1\,080\,000} = 102,6\text{cm}$ $\therefore h = \frac{2\,160\,000}{(102,6)^2} = 205,19\text{cm}$	✓ $8x - 8\,640\,000x^{-2}$ ✓ equating $A'(x)$ to 0. ✓ simplification ✓ 102,6cm ✓ 205,19cm	(5)
			[10]

TOTAL: 150